


EXPLANATORY DATA
SAN LUIS OBISPO SHEET
GEOLOGIC MAP OF CALIFORNIA

OLAF P. JENKINS EDITION
Compiled by Charles W. Jennings, 1958

INDEX TO GEOLOGIC MAPPING
USED IN COMPILATION OF THE SAN LUIS OBISPO SHEET



1. Anonymous, Western portion of Pismo basin, San Luis Obispo County, scale 1" = 2,000 feet, unpublished (1954).
2. Bailey, Edgar H., 1942, Quicksilver deposits of the Parkfield district, California: U. S. Geol. Survey Bull. 936-F, pp. 143-169, Pl. 19: Geologic map and section of the Table Mountain area, scale 1:31,680.
3. Brooks, T. J., 1955, Spring field trip Devils Den—McLure Valley area, California: S.E.P.M.—A.A.P.G., 4 pp., Map: Geologic map Pyramid Hills—Devils Den area, scale 1" = 4,000 feet.
- 4a. Dibblee, Thomas W., Jr., Geologic map of the Cholame quadrangle, California, Scale 1:125,000, unpublished.
- 4b. Dibblee, Thomas W., Jr., Geologic maps of the Pozo, La Panza, and part of Branch Mtn. quadrangles, scale 1:62,500, unpublished.
5. Fairbanks, H. W., 1904, San Luis Folio, California: U. S. Geol. Survey Geologic Atlas (No. 101), 14 pp., Map: Areal Geology sheet, scale 1:125,000.
(Faults interpreted from Fairbanks cross-sections and data from J. O. Kistler, and L. A. Tarbet, Standard Oil Company of California. Quaternary geology in Morro Bay area and Nipomo area modified by C. W. Jennings, California Division of Mines.)
- *6. Heikkila, Henry H., and MacLeod, George M., 1951, Geology of Bitterwater Creek area, Kern County, California: California Division Mines Special Rept. 6, 21 pp., Pl. 1: Topographic and geologic map of the Bitterwater Creek area, Kern County, California, scale 1:31,680.
- *7. Herrera, Leo J., Jr., Geology of the Tent Hills quadrangle, scale 1:31,680, University of California, Berkeley, unpublished Masters thesis, 1950.
- *8. Herron, Robert F., Geologic map of Highland-San Juan area, scale 1:62,500, unpublished (1948).
9. Marsh, Owen T., Geology of the Orchard Peak area, scale 1:31,680, Stanford University, unpublished Ph. D. thesis, 1954.
10. Peryam, Richard C., Geology of the Annette quadrangle, San Luis Obispo and Kern Counties, California, scale 1:31,680, University of California, Berkeley, unpublished M.A. thesis, 1949.
- *11. Stewart, Ralph, 1946, Geology of Reef Ridge, Coalinga district, California: U. S. Geol. Survey Prof. Paper 205-C, pp. 81-115, Pl. 9: Geologic map and sections of Reef Ridge and parts of adjoining regions, Coalinga district, California, scale 1:31,680.
- *12a. Taliaferro, N. L., (and summer field classes), Geologic maps of the Adelaida, Bradley, Bryson, Cape San Martin, Paso Robles, Piedras Blancas, San Miguel, and San Simeon quadrangles, scale 1:62,500, University of California, Berkeley, (compiled in 1957 by R. E. Turner), unpublished.
(Faults in San Antonio River area from L. A. Tarbet, Standard Oil Company of California.)
- Taliaferro, N. L., 1943, Bradley-San Miguel district: California Div. Mines Bull. 118, pp. 456-462, Fig. 189: Bradley and San Miguel quadrangles, geologic map, scale approx. 5/16" = 1 mile.
- Taliaferro, N. L., 1944, Cretaceous and Paleocene of Santa Lucia Range, California: Amer. Assoc. Petroleum Geologists Bull., vol. 28, pp. 449-521, Fig. 9: Map of parts of Adelaida, Bradley, Bryson and San Simeon quadrangles, scale approx. 1-1/16" = 1 mile.
- 12b. Taliaferro, N. L. (and summer field classes), Geologic map of northern part of Parkfield quadrangle, scale 1:31,250, University of California, Berkeley, unpublished.
- *12c. Taliaferro, N. L. (and summer field classes), Geologic maps of Nipomo and part of Branch Mtn. quadrangle, scale 1:62,500: University of California, Berkeley, unpublished. (Modified in part by J. O. Kistler, Standard Oil Company of California.)
- Taliaferro, N. L., 1943, Geology of Huasna area: California Div. Mines Bull. 118, pp. 443-447, Fig. 185: Huasna area, geologic map, scale approx. 1/2" = 1 mi.
13. Woodring, W. P., Stewart, Ralph, and Richards, R. W., 1940, Geology of the Kettleman Hills oil field, California, stratigraphy, paleontology, and structure: U. S. Geol. Survey Prof. Paper 195, 170 pp., Pl. 3: Geologic map of the Kettleman Hills, California, scale 1:31,680.
14. Worts, G. F., Jr., 1951, Geology and ground-water resources of the Santa Maria Valley area, California: U. S. Geol. Survey Water-Supply Paper 1000, 169 pp., Pl. 1: Geologic map of the Santa Maria Valley area, Santa Barbara County, California, scale 1:63,360.

For a complete list of published geologic maps of this area see California Div. Mines Special Report 52.



Aerial view northwestward along axis of McLure Valley syncline. Pliocene Jacalitos formation in center middleground flanked by upper Miocene McLure shale and Upper Cretaceous Panoche sandstone and shale. Photo by John S. Shelton.

STRATIGRAPHIC NOMENCLATURE — SAN LUIS OBISPO SHEET

LEGEND SAN LUIS OBISPO SHEET

DATA FROM SELECTED PUBLISHED SOURCES USED TO COMPILE THE SAN LUIS OBISPO SHEET Numbers Refer to Index on Reverse Side of Sheet

AGE	STATE MAP SYMBOL	STATE MAP UNIT	Brooks 3	Fairbanks 5	Heikkila & MacLeod 6	Stewart 11	Taliaferro 12a (In part unpublished)	Taliaferro 12c	Woodring, Stewart & Richards 13	Worts 14	
CENOZOIC	Recent	Qs	RECENT SAND DUNES (Dune sand)							Dune sand	
		Qal	RECENT ALLUVIUM (Alluvium, river channel deposits)			Alluvium and stream gravel		Alluvium	Alluvium	River-channel deposits, alluvium	
		Qf	RECENT ALLUVIAL FAN DEPOSITS IN THE GREAT VALLEY (Sediments deposited from streams emerging from high lands surrounding the Great Valley)		Alluvium		Alluvium	Alluvium		Younger alluvium	
	Pleistocene	Qt	RIVER AND STREAM TERRACE DEPOSITS (Terrace deposits)			Terrace deposits (and dune sand)	Terrace gravels		Terraces		Terrace deposits
		Qm	PLEISTOCENE MARINE DEPOSITS AND MARINE TERRACE DEPOSITS (Marine terrace deposits)					Marine terrace deposits			
		Qc	PLEISTOCENE NONMARINE SEDIMENTARY DEPOSITS (Orcutt fm.)								Orcutt fm.
	Pliocene	QP	PLIOCENE-PLEISTOCENE NONMARINE SEDIMENTARY DEPOSITS (McKittrick fm., Paso Robles fm., Tulare fm.)		Tulare fm.	Paso Robles fm.	Tulare fm.	Tulare fm.	Paso Robles		Tulare fm.
		Pc	UNDIVIDED PLIOCENE NONMARINE SEDIMENTARY ROCKS (Morales fm., unnamed Pliocene nonmarine beds)								
		Pu	UPPER PLIOCENE MARINE SEDIMENTARY ROCKS (San Joaquin fm.)		San Joaquin Clay			San Joaquin fm.			San Joaquin fm.
	Miocene	Pml	MIDDLE AND LOWER PLIOCENE MARINE SEDIMENTARY ROCKS (Etchegoin fm., Etchegoin-Jacalitos fm. undifferentiated, Jacalitos fm., Pancho Rico fm. [in part upper Miocene])		Etchegoin fm.			Etchegoin fm., Jacalitos fm.	Etchegoin- Jacalitos fm. undifferentiated	Pliocene, chiefly sands and silts	Etchegoin
		Mc	UNDIVIDED MIOCENE NONMARINE SEDIMENTARY ROCKS (Unnamed Miocene continental rocks)						Unnamed Miocene continental red-beds and volcanic rocks		
		Mu	UPPER MIOCENE MARINE SEDIMENTARY ROCKS (Antelope shale, member of Monterey fm., McDonald shale, McLure shale member of Monterey fm., Pismo fm., Reef Ridge shale, Santa Margarita fm.)		McLure shale, McDonald shale	Pismo fm., Santa Margarita fm.	McDonald shale	McLure shale member of Monterey shale and Reef Ridge shale	Santa Margarita, McLure shale	Santa Margarita	
		Mm	MIDDLE MIOCENE MARINE SEDIMENTARY ROCKS (Alferitz fm., "Buttonbed" sandstone, Escuda sand- stone, Gould-Devilwater shale, Monterey shale [in part upper Miocene], Point Sal fm., Salinas shale, Twissle- man sandstone member of Monterey fm.)		Escuda sand	Monterey shale	Twisselmann sandstone mem- ber of Monterey fm., Gould- Devilwater shale, "Buttonbed" sandstone		Monterey fm., Salinas shale	Shale (lower & middle Miocene)	
		MI	LOWER MIOCENE MARINE SEDIMENTARY ROCKS (Agua sandstone, Carneros sandstone, Hannah fm., Media shale, Rincon fm., Sandholt fm., Santos shale, Temblor fm. [middle Miocene in part], Vaqueros sand- stone)		Temblor- Vaqueros fm.	Vaquero sandstone	Media shale, Carneros ss, Upper Santos shale, Agua sandstone	Temblor sandstone	Vaqueros	Vaqueros sandstone	
		Mv ^a Mv ^b Mv ^P	MIOCENE VOLCANIC ROCKS: UNDIFFERENTIATED—Mv; ANDESITIC —Mv ^a ; BASALTIC—Mv ^b ; PYROCLASTIC —Mv ^P (Undifferentiated volcanic rocks, teschenite—Mv; andesite—Mv ^a ; diabase—Mv ^b ; Obispo tuff, rhyolite tuff—Mv ^P)			Augite- teschenite—Mv, pyroxene andesite—Mv ^a , olivine diabase— Mv ^b , rhyolite tuff—Mv ^P			Undifferentiated rhyolite tuff and diabase sills interbedded in Monterey fm.— Mv		
	Oligocene	Φc	OLIGOCENE NONMARINE SEDIMENTARY ROCKS (Berry conglomerate)								
		Eocene	E	EOCENE MARINE SEDIMENTARY ROCKS (Avenal sandstone, Canoas siltstone, Gredal fm., Krey- enhagen shale, Mabury fm., Point of Rocks sandstone, Welcome fm.)		Kreyenhagen shale, Point of Rocks sandstone, Canoas siltstone, Avenal sandstone		Kreyenhagen shale, Point of Rocks sandstone	Kreyenhagen shale, Avenal sandstone		
	Ep		PALEOCENE MARINE SEDIMENTARY ROCKS (Dip Creek fm.)						Dip Creek fm.		
Undivided	Ti	TERTIARY INTRUSIVE ROCKS (Plugs and dikes of quartz porphyry, andesite and dacite.)			Andesite- granophyre ² dacite- granophyre ²			Middle Miocene intrusive dacite	Quartz porphyry plugs and dikes		

STRATIGRAPHIC NOMENCLATURE — Continued

AGE	STATE MAP SYMBOL	STATE MAP UNIT	Brooks 3	Fairbanks 5	Heikkila & MacLeod 6	Stewart 11	Taliaferro 12a (In part unpublished)	Taliaferro 12c	Woodring, Stewart & Richards 13	Worts 14
MESOZOIC { CRETACEOUS { JURASSIC { PALEOZOIC { UNDIVIDED	K	UNDIVIDED CRETACEOUS MARINE SEDIMENTARY ROCKS (Unnamed undivided Cretaceous of Table Mt. area)			Cretaceous undifferentiated			Cretaceous undifferentiated		
	Ku	UPPER CRETACEOUS MARINE SEDIMENTARY ROCKS (Asuncion group, Atascadero fm., Jack Creek fm., Moreno fm., Panoche fm.)	Moreno fm.	Atascadero fm.		Panoche fm.	Asuncion group, Jack Creek fm.			
	Kl	LOWER CRETACEOUS MARINE SEDIMENTARY ROCKS (Marmolejo fm., Toro fm.)		Toro fm.			Marmolejo fm.			
	Kjf	FRANCISCAN GROUP (Franciscan fm., San Luis fm.)		Franciscan "San Luis" fm. (sandstone, conglomerate, shale and schist)		Franciscan fm. (sandstones, and volcanic and metamorphic rocks)	Franciscan fm. (sandstone, shale, chert, schist)			Franciscan and Knoxville(?) fms. (sandstone, shale, chert, serpentine and schist)
	Kjfv	FRANCISCAN VOLCANIC AND METAVOLCANIC ROCKS (Cuesta diabase, Osos basalt, altered volcanic rocks)		Cuesta diabase, Osos basalt, diabase and other basic rocks			Franciscan volcanic rocks			
	gr	MESOZOIC GRANITIC ROCKS (Santa Lucia quartz diorite, granite)		Granite			Granitic rocks (considered by N. L. T. to be pre-Franciscan, probably Precambrian)			
	ub	MESOZOIC ULTRABASIC INTRUSIVE ROCKS (Serpentine, peridotite, pyroxenite, norite, gabbro)		Serpentine and related rocks			Serpentine and related ultrabasic rocks			
	Jk	KNOXVILLE FORMATION (Knoxville fm.)					Knoxville fm.			
m ls	PRE-CRETACEOUS METAMORPHIC ROCKS, UNDIFFERENTIATED, ls = LIMESTONE (Sur series, complex of metamorphic and granitic rocks—m; Gabilan limestone and dolomite lenses—ls)					Sur series—m; Gabilan limestone—ls				

NOTES

¹ Etchegoin fm.: mostly middle Pliocene; locally grades upward into overlying nonmarine Paso Robles formation and contains upper Pliocene beds.

² Considered to be of Cretaceous age by Fairbanks; correlated by Taliaferro with Miocene plugs and dikes to north.



View southward to Morro Bay on the San Luis Obispo coast. Morro Rock, a seastack on the extreme right, is one of a group of aligned volcanic plugs appearing as dark hills in the middleground. Beyond Morro Rock, a baymouth bar separates Morro Bay from the open Pacific Ocean. Franciscan formation and associated serpentine intrusive rocks are exposed in the foreground. The dissected San Luis Range in the distance is composed largely of the Monterey formation. *Photo by John S. Shelton.*

TOPOGRAPHIC QUADRANGLES
 WITHIN THE SAN LUIS OBISPO SHEET
 AVAILABLE FROM THE U. S. GEOLOGICAL SURVEY

1958



View south along crest of Reef Ridge toward Pyramid Hills and San Joaquin Valley. Resistant Temblor sandstone forms crest of dark ridge in foreground (A), while light ridge in middleground (B), is composed of unconformably overlying Monterey shale. Younger Pliocene formations extend toward Kettleman Plain on left. *Photo by John S. Shelton.*